



Suggested Uses:

ER:

- Patients without undergarments.
- Patients who need a procedure where their garments are at risk for soiling.
- Patients who are being admitted directly to the OR or to interventional radiology, regardless of procedure. (Exclude procedures that will require Under Johnnies removal i.e appendectomy, ectopic pregnancy, D&C, etc.)
- Patients admitted to the floor where repeated abdominal exams are required.

Radiology:

- Chest/abdominal/rectal interventional procedures (Exclude procedures where access to the groin is necessary . In these cases, use post procedure.)

Preoperatively:

- All surgical procedures that do not require a foley catheter or access to the lower abdomen, perineum or anal area. For example: thyroid surgery, breast surgery, plastic surgery above the umbilicus, thoracic surgery, bariatric surgery, cholecystectomy, gastric tube, jejunostomy tube, ventral hernias, umbilical hernias, orthopedic surgery of the knee, podiatry surgery, etc.

Postoperatively: On transfer off the operating room table.

- All post op patients with surgical drains or foley catheters who did not receive Under Johnnies preoperatively. For example: colectomies, colostomies, prostatectomies, hysterectomies, abdominoplasties, hip surgeries, spine surgery, aortas, vascular surgery.

For Trial Contact: Matthew Traub, OBP Medical, Inc. 1 888 257 9253
matthew.traub@obpmed.com

OBP is proud to provide the medical community with high quality medical devices and innovative solutions to address existing and emerging needs within the medical market. OBP is ISO 13485 certified to ensure production of only the finest in medical garments and equipment.